

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

MS/20190823

1/1

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AP21397BT-N Polyclonal Antibody to Uricase - Biotin

Alternate names:	UOX, Urate oxidase
Quantity:	1 ml
Concentration:	10,0 mg/ml
Uniprot ID:	<u>P16164</u>
NCBI:	<u>NP_999435</u>
GenelD:	<u>397510</u>
Host / Isotype:	Rabbit / IgG
Immunogen:	Uricase isolated and purified from porcine liver. Freund's complete adjuvant is used in the first step of the immunization procedure.
Format:	 State: Lyophilized IgG fraction Purification: Ammonium Sulphate Precipitation and Ion Exchange Chromatography Buffer System: PBS, pH 7.2 without preservatives and foreign proteins Label: Biotin – Marker: N-Hydroxysuccinimidobiotin. Conjugation procedure: A proprietary technique for the binding to biotin is used, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilised to assure stability and long shelf life <i>Molar Ratio:</i> Biotin / IgG ~6.0 Reconstitution: Restore by adding 1.0 ml of sterile distilled water
Applications:	 This product is intended for use in precipitating and non-precipitating antibody- binding assays (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques). <u>Recommended working dilutions:</u> vary widely, but may be between 1/1000 and 1/20000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and enzyme inhibition.
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. If a slight precipitation occurs upon storage, this should be removed by centrifugation. I Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.