

AP21455HR-N**Polyclonal Antibody to Guinea Pig IgM (Fc specific) - HRP**

Alternate names:	Guinea Pig Immunoglobulin M
Quantity:	1 ml
Concentration:	8.0 mg/ml
Host / Isotype:	Goat / IgG
Immunogen:	Purified normal IgM isolated from pooled Guinea Pig serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Format:	State: Lyophilized purified hyperimmune IgG fraction Purification: Ammonium Sulphate Precipitation and Ion Exchange Chromatography Buffer System: PBS, pH 7.2 without preservatives and foreign proteins Label: HRP – Horseradish Peroxidase <i>Molar Ratio:</i> Peroxidase/IgG ~ 1.7 Reconstitution: Restore by adding 1.0 ml of sterile distilled water
Applications:	Can be used for Enzyme-Immunocytochemical and immunohistochemical detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgM antibodies in serodiagnostic microbiology and autoimmune diseases. To identify a specific antigen using a reference antibody of Guinea Pig origin known to be of the IgM isotype in the middle layer of the indirect test procedure. In non-isotopic assay methodology (e.g. ELISA) to measure IgM in Guinea Pig serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <i>Recommended Working Dilutions:</i> Histochemical and Cytochemical: 1/100-1/250. ELISA and comparable non-precipitating antibody-binding assays: 1/400-1/4000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in Guinea Pig serum, as tested by Immunoelectrophoresis and Double Radial Immunodiffusion. Cross-Reactivity: Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail.
Add. Information:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

Storage:

Prior to reconstitution store at 2-8°C.
Following reconstitution store undiluted at 2-8°C for one week
or (in aliquots) at -20°C for longer.
Avoid repeated freezing and thawing.
Shelf life: one year from despatch.