

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AP23427PU-S Polyclonal Antibody to HA Epitope Tag (YPYDVPDYA) - Aff -

Purified

Alternate names: HA Tag, HA-Tag, Hemagglutinin Tag

Quantity: 50 μg Concentration: 1.0 mg/ml

Background: Plasmid vectors for the expression of coding regions of eukaryotic genes in acterial,

insect and mammalian hosts are in common usage; such expression ectors are frequently used to encode hybrid fusion proteins consisting of a eukaryotic target protein and a specialized region designed to aid in the urification and visualization of the target protein. For example, the pCDM8 expression vector and derivatives thereof encode fusions between the target protein and an 11 amino acid peptide derived from

the influenza protein hemagglutinin (HA).

Host / Isotype: Rabbit / IgG

Immunogen: A synthetic peptide from influenza hemagglutinin epitope coupled to KLH

AA Sequence: YPYDVPDYA

Format: State: Liquid Ig fraction

Purification: Affinity chromatography

Buffer System: PBS, pH 7.4, with 0.05% sodium azide

Applications: Western blot: 1:500~1:1000.

Immunoflourescence: 1:200-400. Immunoprecipitation: 1:200-400.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: HA-Tag polyclonal antibody detects HA-tagged proteins overexpressed in cells. The

antibody recognizes the HA-tag fused to either the amino- or carboxy-terminus of

targeted proteins in transfected mammalian cells.

Species Reactivity: Tested: Mammalia

Storage: Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: Field, J. et al. (1988) Mol. Cell. Biol. 8, 2159–2165.