

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Schillerstr. 5

AP30526PU-N Polyclonal Antibody to LZTR1 - Aff - Purified

Alternate names: Leucine zipper-like transcription regulator 1

Quantity: 0.1 mg
Concentration: 1.0 mg/ml

Background: LZTR1, a member of the BTB-kelch superfamily, was initially described as a putative

transcriptional regulator based on weak homology to members of the basic leucine zipper-like family, the encoded protein subsequently has been shown to localize exclusively to the Golgi network where it may help stabilize the Golgi complex. Deletion of this gene may be associated with DiGeorge syndrome, a developmental field defect involving the third and fourth pharyngeal pouches, causing the absence of thymus and parathyroid glands, congenital cardiac abnormalities and facial dysmorphism. LZTR1 is tyrosine phosphorylated and subsequently degraded upon

induction of apoptosis.

 Uniprot ID:
 Q20WK0

 NCBI:
 CAI86451

 GeneID:
 8216

Host / Isotype: Rabbit / IgG

Immunogen: LZTR1 antibody was raised against a 14 amino acid peptide near the amino terminus

of human LZTR1.

Format: State: Liquid Ig fraction

Purification: Peptide affinity chromatography **Buffer System:** PBS containing 0.02% sodium azide

Applications: ELISA.

Western blot: 1 - 2 μg/ml.

Immunohistochemistry on paraffin sections.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Species Reactivity: Tested: Human, mouse, rat

Add. Information: Blocking peptide available: AP30526CP-N

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Shelf life: one year from despatch.

General Readings: 1. Kurahashi H, Akagi K, Inazawa J, Ohta T, Niikawa N, Kayatani F, et al. Isolation and

characterization of a novel gene deleted in DiGeorge syndrome. Hum Mol Genet. 1995

Apr;4(4):541-9. PubMed PMID: 7633402.


2. Nacak TG, Leptien K, Fellner D, Augustin HG, Kroll J. The BTB-kelch protein LZTR-1 is a novel Golgi protein that is degraded upon induction of apoptosis. J Biol Chem. 2006

Feb 24;281(8):5065-71. Epub 2005 Dec 15. PubMed PMID: 16356934.



Pictures:

Western blot analysis of LZTR1 in human heart tissue lysate with LZTR1 antibody at (A) 1 and (B) 2 ug/mL.



Immunohistochemistry of LZTR1 in mouse heart tissue with LZTR1 antibody at 5 ug/mL.

