

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AP31483FC-N Polyclonal Antibody to Mouse Serum Proteins - FITC

Quantity: 1 ml

Concentration: 10 mg/ml
Host: Goat

Immunogen: Pooled whole serum and selected protein fractions isolated from pooled mouse

serum. Freund's complete adjuvant is used in the first step of the immunization

procedure.

Format: State: Lyophilised hyperimmune Ig fraction

Purification: DEAE-column Chromatography

Buffer System: PBS, pH 7.2

No preservative added. No foreign proteins added. **Label:** FITC – Fluorescein isothiocyanate isomer 1

Absorption / Emission: 492nm / 515nm

Molar Ratio: 1,5

Reconstitution: Restore with 1 ml sterile distilled water

Applications: Direct fluorescent staining of mouse serum proteins at the cellular and subcellular

level of appropriately treated cell and tissue substrates. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Working dilutions are usually between 1/20 and 1/80.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: Tested in immunoelectrophoresis and double radial immunodiffusion against pooled

serum precipitation of at least 15 individual proteins have been observed.

<u>Cross-reactivity:</u> Inter-species cross-reactivity is a normal feature of antibodies to serum proteins, since homologous proteins of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in

detail.

Species: Mouse.

Other species not tested.

Storage: Prior to reconstitution store at 2-8°C.

Following reconstitution store undiluted at 2-8°C for one week

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.