

**AP31540BT-N****Polyclonal Antibody to Canine IgG (Fc specific) - Biotin**

<b>Alternate names:</b>	dog IgG, dog immunoglobulin G
<b>Quantity:</b>	1 ml
<b>Concentration:</b>	10 mg/ml
<b>Host / Isotype:</b>	Goat / IgG
<b>Immunogen:</b>	Purified normal IgG isolated from pooled dog serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Format:</b>	<b>State:</b> Lyophilised hyperimmune Ig fraction <b>Purification:</b> DEAE-column Chromatography <b>Buffer System:</b> PBS, pH 7.2 No preservative added, as it may interfere with the antibody activity. No foreign proteins added. <b>Label:</b> Biotin <i>Molar Ratio: 6,7</i> <b>Reconstitution:</b> Restore with 1 ml sterile distilled water
<b>Applications:</b>	Can be used in immunocytochemical and immunohistochemical use for the detection of IgG at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgG antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using an reference antibody of goat origin known to be of the IgG isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgG in dog serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. <b>Working dilutions:</b> For <b>histochemical and cytochemical</b> use are usually between 1/100 and 1/250. In <b>ELISA and comparable non-precipitating antibody-binding assays</b> between 1/500 and 1/5000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in dog serum, as tested by immunoelectrophoresis and double radial immunodiffusion. <b>Cross-reactivity:</b> Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiserum has not been tested in detail. <b>Species:</b> Canine. Other species not tested.

**Add. Information:**

**Adsorption:** Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

**Storage:**

Prior to reconstitution store at 2-8°C.  
Following reconstitution store undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer.  
Avoid repeated freezing and thawing.  
Shelf life: one year from despatch.