

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Schillerstr. 5

AP33041SU-N Polyclonal Antibody to Bisphenol A - Serum

Quantity: 0.1 ml

Background: Bisphenol A (BPA) is predominantly used in the production of polycarbonate plastics

and epoxy resins that are used in many products. It is released into the environment and food. BPA is an endocrine disruptor with estrogenic and obesogenic properties. It influences reproduction and has an epigenetic effect already in the foetus. Bisphenol A is a known hormone-disrupting agent, commonly used in plastics and diffusing into

the environment and food.

Host: Rabbit

Immunogen: BSA-Bisphenol Valeric Acid (BVA)

Format: State: Liquid Serum

Applications: ELISA: Dilution of 1/50,000 from the delivered immune serum. Plates are coated with

400 ng/ml OVA-Bisphenol Valeric Acid (BVA). HRP-conjugated anti-Rabbit IgG as a

tracer 1/8,000.

Immunoaffinity Chromatography.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: Target: Bisphenol A, CAS no.: 80-05-7, Solubility: Acetone, Ethanol, Ether, Benzene.

This antibody is highly specific for Bisphenol A.

Cross Reactivity in a Direct Assay

Molecules containing a phenolic group: Bisphenol A: 100%, 4,4'-(ethylidene) bisphenol: 10%, Bis-(4-hydroxy phenyl)-methane: 1%, Nonylphenol: 1%,

4-cumylphenol: 20%.

Molecules lacking a phenolic group: Vinclozolin: 0.1%, Pirimifos-ethyl: < 0.1%, 17ß-

Estradiol: < 0.1%, 2,4 D: 0.1%, Sulfadimidine: < 0.1%.

Cross Reactivity in an Indirect Assay

Molecules containing a phenolic group: Bisphenol A: 100%, 4,4'-(ethylidene) bisphenol: 100%, Bis-(4-hydroxy phenyl)-methane: 4-cumylphenol: 100%.

Molecules lacking a phenolic group: Vinclozolin: < 0.1%, Pirimifos-ethyl: < 0.1%, 2,4 D:

0.1%, Fenitrothion: < 0.1%, Chlorpyrifos-methyl: < 0.1%, Erythromycine: < 0.1%.

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: 1. Eline P. Meulenberg, Kees Koopal and Ria Rhemrev - Immunoassays for

alkylphenolic pollutants with endocrine disrupting activity. Intern. J. Environ. Anal.

Chem. Vol 85, no 12-13, 15 October-15 November 2005, 871-883.