

AP33059SU-N**Polyclonal Antibody to Guinea Pig IgA (Fc specific) - Serum****Alternate names:**

Guinea Pig Immunoglobulin A

Quantity:

1 ml

Concentration:

Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.

Host:

Sheep

Immunogen:Highly purified normal IgA isolated from pooled guinea pig serum.
Freund's complete adjuvant is used in the first step of the immunization procedure.**Format:****State:** Lyophilized (delipidated, heat inactivated) stable whole antiserum without preservatives**Reconstitution:** Restore by adding 1 ml of sterile distilled water**Applications:****Immunoprecipitation.**

Can be used in precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgA in guinea pig serum or other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of guinea pig IgA from serum or plasma. This antiserum is not intended for use in non-precipitating antibodybinding or other highly sensitive assays. This does not exclude such use if [proper controls are included.

Directions for Use:

Immunoelectrophoresis: Use 2 µl or equivalent against 120 µl antiserum.

Double Radial Immunodiffusion use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells.

Antibody titre: Precipitin titre not less than 1/40 when tested against normal guinea pig serum in agar block titration.

Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.

Specificity:

The reactivity of the antiserum is restricted to the Fc part of the IgA molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal guinea pig serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgA. No precipitation reaction is obtained with purified IgG, IgM, and IgG/Fab fragments.

Cross-reactivity: The antiserum does not cross-react with any other component of the guinea pig immunoglobulin system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Crossreactivity of this antiserum has not been tested in detail.**Species:** Guinea Pig.

Other species not tested.

Add. Information:

Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required, to eliminate antibodies cross-reacting with other components of the immunoglobulin system or reacting with other serum proteins. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.

Storage:

Store lyophilized at 2-8°C for 6 months or at -20°C long term.
After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term.
Avoid repeated freezing and thawing.
Shelf life: one year from despatch.