


## Polyclonal Antibody to C17orf65 (N-term) - Aff - Purified

<b>Alternate names:</b>	Uncharacterized protein
<b>Catalog No.:</b>	AP50499PU-N
<b>Quantity:</b>	0.1 mg
<b>Concentration:</b>	0.25 mg/ml
<b>Uniprot ID:</b>	<a href="#">Q495Z4</a>
<b>NCBI:</b>	<a href="#">NP_848637</a>
<b>GeneID:</b>	<a href="#">339201</a>
<b>Host / Isotype:</b>	Rabbit / Ig
<b>Immunogen:</b>	KLH conjugated synthetic peptide between 8-37 amino acids from the N-terminal region of human C17orf65
<b>Format:</b>	<b>State:</b> Liquid purified Ig fraction <b>Purification:</b> Affinity chromatography on Protein A <b>Buffer System:</b> PBS containing 0.09% (W/V) sodium azide as preservative
<b>Applications:</b>	<b>ELISA:</b> 1/1000. <b>Western blotting:</b> 1/100 - 1/500. <b>Immunohistochemistry on paraffin sections:</b> 1/50 - 1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody reacts to C17orf65.
<b>Add. Information:</b>	<b>Molecular Weight:</b> 20099 Da
<b>Storage:</b>	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
<b>General Readings:</b>	Wang, A.G., et al. Biochem. Biophys. Res. Commun. 345(3):1022-1032(2006)

**Pictures:**

C17orf65 Antibody (N-term) (Cat. #AP50499PU-N) immunohistochemistry analysis in formalin fixed and paraffin embedded human bladder carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of C17orf65 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



C17orf65 Antibody (N-term) (Cat. #AP50499PU-N) western blot analysis in MDA-MB231 cell line lysates (35µg/lane). This demonstrates the C17orf65 antibody detected the C17orf65 protein (arrow).

