

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com AP52372PU-N
OriGene EU

Jildelle EU

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Polyclonal Antibody to KIR2DS5 (C-term) - Aff - Purified

Alternate names: CD158 antigen-like family member G, CD158G, CD_antigen=CD158g, Flags: Precursor, Killer

cell immunoglobulin-like receptor 2DS5, MHC class I NK cell receptor, NKAT-9, NKAT9,

Natural killer-associated transcript 9

Catalog No.: AP52372PU-N

Quantity: 0.4 ml
Concentration: lot specific

Background: Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins

expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several 'framework' genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are

thought to play an important role in regulation of the immune response.

 Uniprot ID:
 Q7L8K3

 NCBI:
 NP_055328

GeneID: 3810
Host / Isotype: Rabbit / Ig

Immunogen: KLH conjugated synthetic peptide selected from the C-terminal region of Human KIR2DS5.

Genename: KIR2DS5

Format: State: Liquid purified Ig fraction

Purification: Protein A column, followed by peptide affinity purification

Buffer System: PBS

Preservatives: 0.09% (W/V) Sodium Azide

Applications: ELISA: 1/1000.

Western Blot: 1/100-1/500.


Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognizes Human KIR2DS5 (C-Term).

Add. Information: Molecular Weight: 33644 Da







Pictures:

Western blot analysis of KIR2DS5 Antibody (C-Term) Cat.-No AP52372PU-N in CEM cell line lysates (35ug/lane). KIR2DS5 (arrow) was detected using the purified Pab.

