



## Polyclonal Antibody to MAGEB17 (N-term) - Aff - Purified

<b>Alternate names:</b>	Melanoma-associated antigen B17
<b>Catalog No.:</b>	AP52587PU-N
<b>Quantity:</b>	0.1 mg
<b>Concentration:</b>	0.25 mg/ml
<b>Uniprot ID:</b>	<a href="#">A8MXT2</a>
<b>NCBI:</b>	<a href="#">XP_001130425</a>
<b>GeneID:</b>	<a href="#">645864</a>
<b>Host / Isotype:</b>	Rabbit / Ig
<b>Immunogen:</b>	KLH conjugated synthetic peptide between 84-114 amino acids from the N-terminal region of human MAGEB17
<b>Format:</b>	<b>State:</b> Liquid purified Ig fraction <b>Purification:</b> Protein A column, followed by peptide affinity purification <b>Buffer System:</b> PBS containing 0.09% (W/V) Sodium Azide as preservative
<b>Applications:</b>	<b>ELISA:</b> 1/1000. <b>Western Blot:</b> 1/100-1/500. <b>Immunofluorescence:</b> 1/10-1/50. <b>Immunohistochemistry on Paraffin Sections:</b> 1/10-1/50. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognizes Human MAGEB17 (N-term).
<b>Add. Information:</b>	<b>Molecular Weight:</b> 51128 Da
<b>Storage:</b>	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
<b>Pictures:</b>	Immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue reacted with MAGEB17 Antibody (N-term) Cat.-No AP52587PU-N, which was peroxidase conjugated to the secondary antibody and followed by AB staining. 

Confocal immunofluorescent analysis of MAGEB17 Antibody (N-term)  
Cat.-No AP52587PU-N with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Western blot analysis of MAGEB17 Antibody (N-term) Cat.-No AP52587PU-N in HeLa,HL-60 cell line lysates (35ug/lane). This demonstrates the MAGEB17 antibody detected the MAGEB17 protein (arrow).

