



AP54734PU-N**Polyclonal Antibody to ZNF98 (C-term) - Aff - Purified**

Alternate names:	ZNF739, Zinc finger protein 98
Quantity:	0.4 ml
Concentration:	lot specific
Background:	May be involved in transcriptional regulation.
Uniprot ID:	A6NK75
NCBI:	NP_001092096
GeneID:	148198
Host / Isotype:	Rabbit / Ig
Immunogen:	KLH conjugated synthetic peptide between 486-516 amino acids from the C-terminal region of human ZNF98
Format:	State: Liquid Ig fraction Purification: Protein A column followed by peptide affinity purification Buffer System: PBS with 0.09% (W/V) sodium azide
Applications:	ELISA: 1:1;000. Western blot: 1:100~500. Immunohistochemistry: 1:50~100. Flow cytometry: 1:10~50. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody detects ZNF98 (C-term).
Species Reactivity:	Tested: Human
Add. Information:	Molecular Weight: 61144 Da
Storage:	Store at 2 - 8 °C for up to six months or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
Pictures:	ZNF98 antibody (C-term) (AP54734PU-N) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphnode followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF98 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



ZNF98 Antibody (C-term) (AP54734PU-N) western blot analysis in MDA-MB435 cell line lysates (35 µg/lane). This demonstrates the ZNF98 antibody detected the ZNF98 protein (arrow).



ZNF98 Antibody (C-term) (AP54734PU-N) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

