

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

# AP60003PU-N Polyclonal Antibody to ESM1 - Purified

Alternate names: Endocan, Endothelial cell-specific molecule 1

Quantity: 0.1 mg

Background: Endocan, also known as endothelial cell-specific molecule1 (ESM1), is a secreted

cysteine-rich dermatan sulfate (DS) proteoglycan primarily expressed by endothelial cells within the vascular capillary network in kidney and in the alveolar walls of the lung. Endocan expression has also been detected in different epithelia and in

adipocytes. The expression of endocan is up-regulated by TNFα, IL1β or

lipopolysaccharide and down-regulated by IFNy. The mouse Endocan gene encodes a 184 amino acid (aa) residues precursor protein with a 19 aa hydrophobic signal peptide and a 165 aa mature region with 18 Cysteine residues. The DS chain is covalently attached to serine 137. Endocan has been shown to bind CD11a/CD18 integrin (also known as lymphocyte function-associated antigen1, LFA1) on human lymphocytes, monocytes and Jurkat cells, inhibiting its binding to ICAM1 and reducing LFA1mediated leukocyte activation. Endocan binds via its DS chain to hepatocyte growth factor (HGF) to enhance HGF mitogenic activity. Genetically engineered cells overexpressing Endocan has been shown to induce tumor formation, suggesting that Endocan may be involved in the pathophysiology of tumor growth in vivo. Circulating

Endocan can be detected in the serum from healthy subjects.

Uniprot ID: <u>Q9QYY7</u>

NCBI: NP 076101.1

GenelD: <u>71690</u>

Host / Isotype: Rabbit / IgG

Immunogen: Highly pure (>95%) recombinant human Endocan/ESM1 (Trp19-Arg184) derived from E.

coli

Format: State: Lyophilized purified Ig fraction

**Purification:** Protein A Chromatography **Buffer System:** 5 mM PBS, pH 7.2

Reconstitution: Centrifuge vial prior to opening. Restore in sterile water to a

concentration of 0.1-1.0 mg/ml.

**Applications:** Western blot: 1-5 μg/ml.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

**Specificity:** This antibody detects Mouse Endocan / ESM1.

Other species not tested.



#### **Storage:**

Prior to reconstitution store at 2-8°C.

Following reconstitution store undiluted at 2-8°C for one month

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

### **General Readings:**

1. Lassalle P, Molet S, Janin A, Heyden JV, Tavernier J, Fiers W, et al. ESM-1 is a novel human endothelial cell-specific molecule expressed in lung and regulated by cytokines. J Biol Chem. 1996 Aug 23;271(34):20458-64. PubMed PMID: 8702785. 2. Bechard D, Meignin V, Scherpereel A, Oudin S, Kervoaze G, Bertheau P, et al. Characterization of the secreted form of endothelial-cell-specific molecule 1 by specific monoclonal antibodies. J Vasc Res. 2000 Sep-Oct;37(5):417-25. PubMed PMID: 11025405.

- 3. Wellner M, Herse F, Janke J, Gorzelniak K, Engeli S, Bechart D, et al. Endothelial cell specific molecule-1--a newly identified protein in adipocytes. Horm Metab Res. 2003 Apr;35(4):217-21. PubMed PMID: 12778364.
- 4. Béchard D, Gentina T, Delehedde M, Scherpereel A, Lyon M, Aumercier M, et al. Endocan is a novel chondroitin sulfate/dermatan sulfate proteoglycan that promotes hepatocyte growth factor/scatter factor mitogenic activity. J Biol Chem. 2001 Dec 21;276(51):48341-9. Epub 2001 Oct 5. PubMed PMID: 11590178.
- 5. Béchard D, Scherpereel A, Hammad H, Gentina T, Tsicopoulos A, Aumercier M, et al. Human endothelial-cell specific molecule-1 binds directly to the integrin CD11a/CD18 (LFA-1) and blocks binding to intercellular adhesion molecule-1. J Immunol. 2001 Sep 15;167(6):3099-106. PubMed PMID: 11544294.
- 6. Scherpereel A, Gentina T, Grigoriu B, Sénéchal S, Janin A, Tsicopoulos A, et al. Overexpression of endocan induces tumor formation. Cancer Res. 2003 Sep 15;63(18):6084-9. PubMed PMID: 14522939.

## **Pictures:**

Western analysis using anti-mouse Endocan/ESM1 antibody Cat.-No. AP60003PU-N. Samples were loaded in 15% SDS-polyacrylamide gel under reducing conditions.

Lane 1: MWM (kDa), Lane 2: rm ESM1, Lane 3: rh ESM1.

