

**AR08016FC-N****Feline IgG F(c) Fragment**

<b>Alternate names:</b>	Cat IgG, Cat Immunoglobulin G
<b>Quantity:</b>	1 mg
<b>Concentration:</b>	1.0 mg/ml (by UV absorbance at 280 nm)
<b>Species:</b>	Feline
<b>Source:</b>	Serum
<b>Format:</b>	<b>State:</b> Lyophilized IgG F(c) fragment <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA) and 0.01% (w/v) Sodium Azide <b>Label:</b> Fluorescein isothiocyanate (FITC) (Molecular Weight 390 daltons) <b>Protein Ratio:</b> 5.0 moles FITC per mole of Cat IgG F(c) <b>Absorption/Emission:</b> 495 nm / 528 nm <b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent). For extended storage reconstitute product with 50% glycerol instead of water.
<b>Applications:</b>	Flow Cytometry. Immunofluorescence. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Description:</b>	Fluorescein Conjugated Cat IgG F(c) fragment.
<b>Storage:</b>	Prior to reconstitution store at 2-8°C. Following reconstitution store the antigen undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.