


AR09249PU-L

Human DFFA / ICAD (1-331, His-tag) - Purified

Alternate names:	DFF-45, DFF1, DFF45, DNA fragmentation factor 45 kDa subunit, DNA fragmentation factor subunit alpha, Inhibitor of CAD
Quantity:	0.5 mg
Concentration:	1.0 mg/ml (determined by Bradford assay)
Background:	DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA, also known as DNA fragmentation factor subunit alpha, is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis.
Uniprot ID:	O00273
NCBI:	AAH07721
GeneID:	1676
Species:	Human
Source:	E. coli
Format:	State: Liquid purified protein Purity: >90% by SDS – PAGE Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 10% glycerol, 1 mM DTT
Description:	Recombinant DFFA protein was expressed in E.coli and purified by using conventional chromatography techniques. AA Sequence: MGSSHHHHH SSGLVPRGSH MEVTGDAGVP ESGEIRTLKP CLLRRNYSRE QHGVAASCLE DLRSKACDIL AIDKSLTPVT LVLAEDGTIV DDDYFLCLP SNTKFVALAS NEKWAYNSD GGTAWISQES FDVDETDSGA GLKWKNVARQ LKEDLSSIIL LSEEDLQMLV DAPCSDLAQE LRQSCATVQR LQHTLQQVLD QREEVRQSKQ LLQLYLQALE KEGSLLSKQE ESKAAFGEV DAVDTGISRE TSSDVALASH ILTALREKQA PELSLSSQDL ELVTKEPKA LAVALNWDIK KTETVQEACE WELALRLQQT QSLHSLRSIS ASKASPPGDL QNPKRARQDP T Molecular weight: 38.7 kDa (351 aa)
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

Pictures:



15% SDS-PAGE (3ug)

Recombinant human DFFA, 1-331 aa, His-tagged: 15% SDS-PAGE (3 μ g)

