

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

## AR10428PU-N OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## **Mouse Leptin Triple Antagonist - Purified**

Catalog No.: AR10428PU-N

Quantity: 0.1 mg

**Concentration:** 0.65 mg/ml (prior to lyophil.)

Species: Mouse Source: E. coli

Format: State: White lyophilized (freeze-dried) powder

**Purity:** > 99.0% as determined by both Gel filtration and SDS-PAGE analysis.

**Purification Method:** Proprietary chromatographic techniques.

**Buffer System:** Lyophilized from a concentrated (0.65 mg/ml) solution with 0.003mM

NaHCO<sub>3</sub>

Reconstitution: Restore in sterile water or sterile 0.4% NaHCO<sub>3</sub>adjusted to pH 8-9, not less

than 100  $\mu\text{g}/\text{ml}\text{,}$  which can then be further diluted with other aqueous solutions.

**Description:** Recombinant Mouse Leptin Antagonist Triple Mutant is a single non-glycosilated

polypeptide chain containing 146 amino and additional Ala at N-terminus acids and having a Molecular Mass of  $\sim$  16 kDa, LEP was mutated, resulting in L39A/D40A/F41A mutant. **Protein content:** Protein quantization was carried out by UV spectroscopy at 280 nm using the absorbency value of 0.21 as the extinction coefficient for a 0.1% (1mg/ml) solution at pH 8.0. This value is calculated by the PC GENE computer analysis program of protein

sequences (IntelliGenetics).

AA Sequence:

The sequence of the first five N-terminal amino acids was determined and was

found to be Ala-Val-Pro-Ile-Gln

**Biological Activity:** Capable of inhibiting Leptin-induced proliferation of BAF/3 cells stably transfected with the long form of Human Leptin Receptor. It also inhibits various Leptin

effects in several in vitro bioassays.

**Storage:** Prior to reconstitution store at 2-8°C for one month or desiccated below -18°C.

Following reconstitution (at > 0.1 mg/ml and up to 2 mM and filter sterilization) store undiluted at 2-8°C or at RT for one month making it suitable for long term infusion studies

using osmotic pumps.

At lower concentration addition of a carrier protein (0.1% HSA or BSA) is suggested.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

