

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

AR10477PU-N

OriGene EU

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Lysostaphin - Purified

Catalog No.: AR10477PU-N

 Quantity:
 5 mg

 Uniprot ID:
 P10547

 NCBI:
 1286

 Source:
 E. coli

Format: State: Liquid sterile filtered lyophilized powder without any additives

Purity: >96.5% as determined by RP-HPLC

Reconstitution: Restore in 20mM Sodium Acetate, pH 4.5, which can then be further

diluted to other aqueous solutions.

Description: Lysostaphin, an Endopeptidase specific for the cell wall peptidoglycan of staphylococci, is

an extremely potent anti-staphylococcal agent. Lysostaphin is used as a research and diagnostic tool. Because it lyses staphylococci efficiently, it is widely used when preparing staphylococcal DNA or other cellular components for genetic and biochemical studies and for the preparation of protoplasts for transformation. Preparation and analysis of bacterial

DNA has become a powerful tool used by clinical and other microbiologists in

epidemiological studies aimed at tracing sources of infection or bacterial contamination. **Protein content:** Protein quantitation was carried out by two independent methods:

1. UV spectroscopy at 280 nm using the absorbency value of 2.02 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer

analysis program of protein sequences (IntelliGenetics).

2. Analysis RP-HPLC, using a calibrated solution of Lysostaphin as a Reference Standard. **Biological Activity:** Determined by the decrease in turbidity of a suspension of heat-killed

Staphylococcus aureus at pH8.0, 30°C.

Enzymatic Activity: Determined to be 4,135 units/mg.

Specific Activity: Measured with a defined activity of 4,000 U/mg.

Storage: Prior to reconstitution store at 2-8°C for one month or desiccated below -18°C.

Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for

longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

