

Streptavidin-NC - Purified

Catalog No.:	AR10507PU-S
Quantity:	20 µg
Concentration:	3 mg/ml
Background:	<p>Streptavidin is a tetrameric protein secreted by <i>Streptomyces avidinii</i> which binds firmly to biotin. Streptavidin is widely used in molecular biology through its unique high affinity for the vitamin biotin. The dissociation constant (K_d) of the biotin-streptavidin complex is about ~10⁻¹⁵ mol/L. The strong affinity recognition of biotin and biotinylated molecules has made streptavidin one of the most important components in diagnostics and laboratory kits. The streptavidin/biotin system has one of the biggest free energies of association (K_{assoc} = 10¹⁴). The complexes are also extremely stable over a wide range of temperature and pH.</p>
Source:	<i>E. coli</i>
Format:	<p>State: Liquid sterile solution containing 50% Glycerol Purity: >90.0% as determined by: (a) Analysis by RP-HPLC. (b) Anion-exchange FPLC. (c) Analysis by reducing and non-reducing SDS-PAGE Silver Stained.</p>
Applications:	<p>Calibrators and controls for immunoassays and western blot standards. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.</p>
Description:	<p>Recombinant Streptavidin-NC produced in <i>E. coli</i> is a single, non-glycosylated, polypeptide chain having a Molecular Mass of 24kDa. Recombinant Streptavidin-NC not only binds to Nitrocellulose membrane readily but also preserves the full biotin binding ability.</p>
Storage:	<p>Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.</p>