

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

## AR10537PU-L OriGene EU

## **Acris Antibodies GmbH**

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Hepatitis A Virus / HAV P3C (1643-1743) - Purified

Catalog No.: AR10537PU-L

Quantity: 1 mg

Concentration: 1.0 mg/ml

**Background:** Forty-two antigenic domains were identified across the hepatitis A virus (HAV) polyprotein

by using a set of 237 overlapping 20-mer synthetic peptides spanning the entire HAV polyprotein. Nineteen antigenic domains were found within the structural proteins, and 22 were found within the nonstructural proteins, with 1 domain spanning the junction of VP1 and P2A proteins. Five of these domains were considered immunodominant, as judged by both the breadth and the strength of their immunoreactivity. One domain is located within the VP2 protein at position 57-90 aa. A second domain, located at position 767-842 aa, contains the C-terminal part of the VP1 protein and the entire P2A protein. A third domain, located at position 1403-1456 aa, comprises the C-terminal part of the P2C protein and the N-terminal half of the P3A protein. The fourth domain, located at position 1500-1519 aa, includes almost the entire P3B, and the last domain, located at position 1719-1764 aa, contains the C-terminal region of the P3C protein and the N-terminal region of the P3D protein. Four of the five most immunoreactive domains are derived from small HAV proteins

and/or encompass protein cleavage sites separating different HAV proteins.

Source: E. coli

Format: Purity: >90% pure as determined by 10% PAGE (coomassie staining).

Purification Method: Inclusion bodies.

Buffer System: 10mM CBB, pH 9.6, 0.1% SDS, 50% Glycerol

Applications: HAV P3C antigen in ELISA and Western blots, excellent antigen for detection of HAV with

minimal specificity problems.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Description:** The *E. Coli* derived 40 kDa recombinant protein contains the P3C immunodominant regions,

amino acids 1643-1743.

**Specificity:** Immunoreactive with sera of HAV-infected individuals.

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.