

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com AR10544PU-N

## OriGene EU

**Acris Antibodies GmbH** 

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Hepatitis B Core / HBcAg (1-183) - Purified

Catalog No.: AR10544PU-N

Quantity: 0.5 mg

Background: Hepatitis B is one of a few known non-retroviral viruses which employ reverse

transcriptionas a part of its replication process. (HIV, a completely unrelated virus, also uses reverse transcription, but it is a retrovirus.) HBV invades the cell by binding to surface receptor and become internalized. The viral core particles then migrate to the hepatocyte nucleus and the partially double-stranded, relaxed circular genomes (RC-DNA) are repaired to form a covalently closed circular DNA (cccDNA), which is the template for viral genomic and sub-genomic RNAs by cellular RNA polymerase II. Of these, the pregenomic RNA (pgRNA is selectively packaged into progeny capsids and is then reverse-transcribed into new RC-DNA. The core can either bud into the endoplasmic reticulum to be enveloped or exported from the cell or recycled back into the genome for conversion to cccDNA.

Source: E. coli

Format: State: 7.5mM Phosphate buffer pH 7.2, 75mM NaCl and 50% Glycerol

**Purity:** HBV Core protein is >95% pure as determined by 10% PAGE (coomassie staining).

Applications: HBV Core antigen is stable in ELISA and Western blots, excellent antigen for detection of

HBV with minimal specificity problems.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Description:** The *E.coli* derived 18 kDa recombinant protein contains the HBV core ayw immunodominant

region, amino acids 1-183.

Specificity: Immunoreactive with sera HBV-infected individuals.

Storage: HBV Core rotein is shipped at ambient temperature.

Upon arrival, Store the antigen at -20°C. Avoid repeated freezing and thawing.