

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

AR10735PU-L

OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Toxoplasma gondii p30 - Purified **Catalog No.:** AR10735PU-L **Quantity:** 1 mg **Concentration:** 1.0 mg/ml **Background:** The life cycle of Toxoplasma gondii has two phases. The coccidia like takes place only in members of the Felidae family which makes these animals the parasite's primary host. The asexual part of the life cycle can take place in any warm-blooded animal, like other mammals(including felines) and birds. T. gondii constructing daughter scaffolds within the mother cell. In the intermediate hosts (including felines), the parasite invades cells, forming intracellular so-called parasitophorous vacuoles containing bradyzoites, the slowly replicating form of the parasit. Vacuoles form tissue cysts mainly within the muscles and brain. Since they are within cells, the host's immune system does not detect these cysts. Resistance to antibiotics varies, but the cysts are very difficult to eradicate entirely. Within these vacuoles T. gondii propagates by a series of binary fissions until the infected cell eventually bursts and tachyzoites are released. Tachyzoites are the motile, asexually reproducing form of the parasite. Unlike the bradyzoites, the free tachyzoites are usually efficiently cleared by the host's immune response, although some manage to infect cells and form bradyzoites, thus maintaining the infection. E. coli Source: **Purity:** >95.0% pure as determined by 10.0% PAGE (coomassie staining). Format: Purification Method: Sepharose-Derived Purification. Buffer System: 50 mM Tris-HCl, 1.5 M Urea, 50% Glycerol **Applications:** Antigen in ELISA and Western blots, excellent antigen for detection of Toxoplasma gondii with minimal specificity problems. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. **Description:** Recombinant Toxoplasma gondii P30 E. coli derived recombinant. The protein contains the P30 (SAG1) immunodominant regions, amino acids 45-198. Specificity: Immunoreactive with sera of Toxoplasma gondii-infected individuals. Store undiluted at 2-8°C for one month or (in aliguots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request. Acris Antibodies is now part of the OriGene family. Learn more at www.origene.com



MS/20120904