

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

AR10736PU-S OriGene EU

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Toxoplasma gondii p29 - Purified

Catalog No.: AR10736PU-S

Quantity: 0.1 mg
Concentration: 1.0 mg/ml

Background: The life cycle of Toxoplasma gondii has two phases. The coccidia like takes place only in members of the Felidae family which makes these animals the parasite's primary host. The asexual part of the life cycle can take place in any warm-blooded animal, like other mammals(including felines) and birds. T. gondii constructing daughter scaffolds within the

mammals(including felines) and birds. T. gondii constructing daughter scaffolds within the mother cell. In the intermediate hosts (including felines), the parasite invades cells, forming intracellular so-called parasitophorous vacuoles containing bradyzoites, the slowly replicating form of the parasit. Vacuoles form tissue cysts mainly within the muscles and brain. Since they are within cells, the host's immune system does not detect these cysts. Resistance to antibiotics varies, but the cysts are very difficult to eradicate entirely. Within these vacuoles T. gondii propagates by a series of binary fissions until the infected cell eventually bursts and tachyzoites are released. Tachyzoites are the motile, asexually reproducing form of the parasite. Unlike the bradyzoites, the free tachyzoites are usually efficiently cleared by the host's immune response, although some manage to infect cells

and form bradyzoites, thus maintaining the infection.

Source: E. coli

Format: Purity: >95.0% pure as determined by 10.0% PAGE (coomassie staining).

Purification Method: Sepharose-Derived Purification. **Buffer System:** 50 mM Tris-HCl, 1.5 M Urea, 50% Glycerol

Applications: Antigen in ELISA and Western blots, excellent antigen for detection of *Toxoplasma gondii*

with minimal specificity problems.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Description: The E.coli derived recombinant protein contains the p29 (GRA7) immunodominant regions,

amino acids 24-100.

Specificity: Immunoreactive with sera of *Toxoplasma gondii*-infected individuals.

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

