

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com


OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AR50928PU-N Human CD74 (73-232, His-tag) - Purified

Alternate names:	DHLAG, HLA class II histocompatibility antigen gamma chain, HLA-DR antigens- associated invariant chain, la antigen-associated invariant chain, p33
Quantity:	0.5 mg
Concentration:	0.5 mg/ml (determined by Bradford assay)
Background:	HLA class II histocompatibility antigen gamma chain isoform b, also known as CD74, is a transmembrane protein which, on antigen presenting cells, is the invariant chain that chaperones MHC class II dimers from the endoplasmic reticulum to the cell surface. CD74 is expressed by cells of both T lymphocyte and B lymphocyte lineages. It is upregulated in several cancers and is also expressed by nonimmune cells during inflammation.
Uniprot ID:	<u>P04233</u>
NCBI:	<u>NP_004346</u>
GenelD:	<u>972</u>
Species:	Human
Source:	E. coli
Format:	State: Liquid purified protein Purity: >90% by SDS - PAGE Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 30% glycerol, 1mM DTT
Description:	Recombinant human CD74 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques. AA Sequence: <u>MGSSHHHHHH SSGLVPRGSH MGSQQQGRLD KLTVTSQNLQ LENLRMKLPK PPKPVSKMRM</u> ATPLLMQALP MGALPQGPMQ NATKYGNMTE DHVMHLLQNA DPLKVYPPLK GSFPENLRHL KNTMETIDWK VFESWMHHWL LFEMSRHSLE QKPTDAPPKE SLELEDPSSG LGVTKQDLGP VPM Molecular weight: 20.6 kDa (183aa) confirmed by MALDI-TOF
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	 Wilson KM, Labeta MO, Pawelec G, Fernandez N. Cell-surface expression of human histocompatibility leucocyte antigen (HLA) class II-associated invariant chain (CD74) does not always correlate with cell-surface expression of HLA class II molecules. Immunology. 1993 Jun;79(2):331-5. PubMed PMID: 8344710. Henne C, Schwenk F, Koch N, Möller P. Surface expression of the invariant chain (CD74) is independent of concomitant expression of major histocompatibility complex class II antigens. Immunology. 1995 Feb;84(2):177-82. PubMed PMID: 7750992.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request. **Pictures:**



15% SDS-PAGE (3ug)

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.