

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford **GERMANY** Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Monoclonal Antibody to Macrophage Galactose Specific Lectin BM4010

(MMGL) - Purified

Alternate names: Connective Tissue Macrophage / Histiocyte Marker

Quantity: 0.1 mg

Concentration: 0.2 mg/ml (stock solution)

Background: MMGL is a glutaraldehyde (0.05%) resistant 38kD single chain surface glycoprotein.

Host / Isotype: Rat / IgG2a **Recommended Isotype**

Controls:

SM15P, SM15PX

Clone: ER-MP23

Immunogen: Mouse macrophage cell lines

Format: State: Lyophilised

Purification: Affinity purified IgG

Buffer System: PBS, pH 7.2 with 10 mg/ml BSA as stabilizer and 0.01% Thimerosal as

preservative

Reconstitution: Restore in 0.5 ml distilled water to make stock solution.

Applications: Immunohistochemistry on frozen sections (Acetone Fixation): Approximate working

dilution is 1 µg/ml (1:200 of stock solution) freshly prepared.

Use mouse skin for positive control.

The antibody has been described to work in Flow cytometry.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

ER-MP23 detects mature macrophages in connective tissues which are situated in the **Specificity:**

vicinity of epithelia e.g. salivary glands, capsule of lymph nodes, thymus and various

other organs.

The antibody blocks the mouse macrophage galactose-specific lectin (MMGL).

Species: Mouse.

Other species not tested.

Add. Information: Staining of isolated cells:

> The antigen is found on the surface of connective tissue macrophages and their precursor cells in bone marrow. Macrophage cell lines (J774-1.6, RAW309Cr.1, WR19M.1) corresponding to mature macrophages express high levels of the antigen.

Staining of tissues:

The antigen is abundant on mature macrophages in connective tissues as present in the skin (dermis), salivary glands, capsule of lymph nodes, thymus and various other

organs. The antigen is partly coexpressed with the BM8 antigen.

ER-MP23 positive macrophages may express la antigens.

Storage: Store the unopend vial at 2-8°C for up to one year and the stock solution in aliquots at

-20°C. Avoid repeated freezing and thawing.

Shelf life: one year from despatch.