

Polyclonal Antibody to Glycogen Synthase pSer641/645 - Aff - Purified

Catalog No.:	BP7077
Quantity:	0.1 ml
Host:	Rabbit
Immunogen:	Chemically synthesized phosphopeptide derived from the region of human Glycogen Synthase that contains serine 641 and serine 645. Remarks: The sequence is conserved in human, mouse and rat.
Format:	State: Liquid Ig fraction Purification: Sequential epitope-specific chromatography. The antibody has been negatively preadsorbed using a non-phosphopeptide corresponding to the site of phosphorylation to remove antibody that is reactive with non-phosphorylated Glycogen Synthase. The final product is generated by affinity chromatography using a Glycogen Synthase-derived peptide that is phosphorylated at serine 641 and serine 645. Buffer System: Dulbecco's phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.3 (+/- 0.1), 50% glycerol with 1.0 mg/mL BSA (IgG, protease free) as a carrier, containing 0.05 %sodium azide as preservative
Applications:	Western blot (starting dilution 1:1,000). Previous lots of this antibody have been used in ELISA. Positive control: Human HepG2 lysates; mouse 3T3-L1 adipocytes lysates. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody detects Glycogen Synthase. Species: Human, mouse, rat. Other species not tested.
Storage:	Store at 2 - 8 °C up to one week or (in aliquots) at -20 °C for longer. Centrifuge vial before opening. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	Takadera, T. and T. Ohyashiki (2004) Glycogen synthase kinase-3 inhibitors prevent caspase-dependent apoptosis induced by ethanol in cultured rat cortical neurons. <i>Eur. J. Pharmacol.</i> 499(3):239-245. Nielson, J.N. and J.F. Wojtaszewski (2004) Regulation of glycogen synthase activity and phosphorylation by exercise. <i>Proc. Nutr. Soc.</i> 63(2):233-237. Ugi, S. et al. (2004) Protein Phosphatase 2A negatively regulates insulin's metabolic signaling pathway by inhibiting Akt (Protein Kinase B) activity in 3T3-L1 adipocytes. <i>Mol. Cell. Biol.</i> 24(19):8778-8789. Grenn, E.R. et al. (2004) Protein-targeting-to-glycogen promotes glycogen storage in hepatocytes by inactivation of phosphorylase and enzyme translocation. <i>J. Biol. Chem.</i>

[Epub ahead of print].

Ugi, S. (2004) Protein Phosphatase 2A negatively regulates insulin's metabolic signaling pathway by inhibiting Akt (Protein Kinase B) activity in 3T3-L1 adipocytes. *Mol. Cell. Biol.* 24(19):8778-8789.

Green, A.R., et al. (2004) Protein-targeting-to-glycogen promotes glycogen storage in hepatocytes by inactivation of phosphorylase and enzyme translocation. *J. Biol. Chem.* [Epub ahead of print].

Kuma, Y., et al. (2004) Identification of glycogen synthase as a new substrate for stress-activated protein kinase 2b/p38beta. *Biochem. J.* 379(Pt 1):133-139.

Skurat, A.V. and A.D. Dietrich (2004) Phosphorylation of Ser640 in muscle glycogen synthase by DYRK family protein kinases. *J. Biol. Chem.* 279(4):2490-2498.

Protocols:

Western Blotting Procedure

1. Lyse approximately 10×10^7 cells in 0.5 mL of ice cold Cell Lysis Buffer (formulation provided below). This buffer, a modified RIPA buffer, is suitable for recovery of most proteins, including membrane receptors, cytoskeletal-associated proteins, and soluble proteins. Other cell lysis buffer formulations, such as Laemmli sample buffer and Triton-X 100 buffer, are also compatible with this procedure. Additional optimization of the cell stimulation protocol and cell lysis procedure may be required for each specific application.
2. Remove the cellular debris by centrifuging the lysates at $14,000 \times g$ for 10 minutes. Alternatively, lysates may be ultracentrifuged at $100,000 \times g$ for 30 minutes for greater clarification.
3. Carefully decant the clarified cell lysates into clean tubes and determine the protein concentration using a suitable method, such as the Bradford assay. Polypropylene tubes are recommended for storing cell lysates.
4. React an aliquot of the lysate with an equal volume of 2x Laemmli Sample Buffer (125 mM Tris, pH 6.8, 10% glycerol, 10% SDS, 0.006% bromophenol blue, and 130 mM dithiothreitol [DTT]) and boil the mixture for 90 seconds at 100°C .
5. Load 10-30 μg of the cell lysate into the wells of an appropriate single percentage or gradient minigel and resolve the proteins by SDS-PAGE.
6. In preparation for the Western transfer, cut a piece of PVDF membrane slightly larger than the gel. Soak the membrane in methanol for 1 minute, then rinse with ddH₂O for 5 minutes. Alternatively, nitrocellulose may be used.
7. Soak the membrane, 2 pieces of Whatman paper, and Western apparatus sponges in transfer buffer (formulation provided below) for 2 minutes.
8. Assemble the gel and membrane into the sandwich apparatus.
9. Transfer the proteins at 140 mA for 60-90 minutes at room temperature.
10. Following the transfer, rinse the membrane with Tris buffered saline for 2 minutes.
11. Block the membrane with blocking buffer (formulation provided below) for one hour at room temperature or overnight at 4°C .
12. Incubate the blocked blot with primary antibody at a 1:1,000 starting dilution in Tris buffered saline supplemented with 3% Ig-free BSA and 0.1% Tween 20 overnight at 4°C or for two hours at room temperature.
13. Wash the blot with several changes of Tris buffered saline supplemented with 0.1% Tween 20.
14. Detect the antibody band using an appropriate secondary antibody, such as goat F(ab)₂ anti-rabbit IgG alkaline phosphatase conjugate or goat F(ab)₂ anti-rabbit IgG horseradish peroxidase conjugate in conjunction with your chemiluminescence reagents and instrumentation.

Cell Lysis Buffer Formulation:

10 mM Tris, pH 7.4

100 mM NaCl
1 mM EDTA
1 mM EGTA
1 mM NaF
20 mM Na₄P₂O₇
2 mM Na₃VO₄
0.1% SDS
0.5% sodium deoxycholate
1% Triton-X 100
10% glycerol
1 mM PMSF (made from a 0.3 M stock in DMSO)
or 1 mM AEBSF (water soluble version of PMSF)
60 µg/mL aprotinin
10 µg/mL leupeptin
1 µg/mL pepstatin
(alternatively, protease inhibitor cocktail such as Sigma Cat. # P2714 may be used)

Transfer Buffer Formulation:

2.4 gm Tris base
14.2 gm glycine
200 mL methanol
Q.S. to 1 liter, then add 1 mL 10% SDS.
Cool to 4°C prior to use.

Tris Buffered Saline Formulation:

20 mM Tris-HCl, pH 7.4
0.9% NaCl

Blocking Buffer Formulation:

100 mL Tris buffered saline
5 gm BSA
0.1 mL Tween 20

Peptide Competition Experiment

The specificity of a Phosphorylation Site Specific Antibody (PSSA) in each experimental system can be confirmed through peptide competition. In this technique, aliquots of antibody are pre-incubated with peptide containing the sequence of the phosphopeptide immunogen used to raise the PSSA and the corresponding non-phosphopeptide. Following preincubation with the peptide, each antibody preparation is then used as a probe in antibody-based detection methods, such as Western blotting, immunocytochemistry, flow cytometry, or ELISA. With a PSSA specific for the phosphorylated target protein, pre-incubation with an excess of peptide containing the sequence of the phosphopeptide immunogen will block all antigen binding sites, while pre-incubation with the corresponding non-phosphopeptide will not affect the antibody.

In performing the Peptide Competition Experiment, it is important to note that the optimal dilutions of both antibody and peptide should be determined empirically for each specific application. The optimal dilution of antibody in these procedures is below saturating, as determined by previous experiments in your system.

The optimal dilution of peptide used in these procedures will depend on the overall affinity or avidity of the antibody, as well as the quantity of the target antigen. A 50-150 fold molar excess of peptide to antibody is found to be effective for most peptide competition

experiments.

In the example presented below, the PSSA is used as a dilution of 1:1000 and the peptides are used at a concentration of 333 nM. The total volume of the phosphopeptide and nonphosphopeptide pre-incubated antibody preparations is 2 mL, sufficient for probing Western blot strips, as well as for use in other antibody-based detection methods. Under these conditions, the molar excess of peptide to antibody is $\times / = 50$.

Procedure:

1. Prepare three identical test samples, such as identical PVDF or nitrocellulose strips to which the protein of interest has been transferred. The test samples should be blocked using a blocking buffer, such as Tris buffered saline supplemented with 0.1% Tween 20, and either 5% BSA or 5% non-fat dried milk.
2. Prepare 6.5 mL of working antibody stock solution (1:1000 in this example) by adding 6.5 μ L of antibody stock solution to 6.5 mL of buffer containing blocking protein, such as TBS supplemented with 0.1% Tween 20, and either 3% BSA or 3% non-fat dried milk.
3. Apportion the unused PSSA into working aliquots and store at -20°C for future use (the stock PSSA contains 50% glycerol and will not freeze at this temperature).
4. Allow the lyophilized control peptides to reach room temperature, ideally under desiccation.
5. Reconstitute each of the control peptides to a concentration of 66.7 μM with nanopure water. (i.e. for a peptide with a molecular mass of 1500, reconstitution with 1 mL water yields a solution with a concentration of 66.7 μM).
6. Apportion the unused reconstituted peptide solutions into working aliquots and store at -20°C for future use.
7. Label 3 test tubes as follows:
 - tube 1: water only no peptide control
 - tube 2: phosphopeptide
 - tube 3: non-phosphopeptide
8. Into each tube, pipette the following components
 - tube 1: 2 mL diluted PSSA solution plus 10 μ L nanopure water
 - tube 2: 2 mL diluted PSSA solution plus 10 μ L phosphopeptide
 - tube 3: 2 mL diluted PSSA solution plus 10 μ L non-phosphopeptide
9. Incubate the three tubes for 30 minutes at room temperature with gentle rocking. During this incubation, the peptides have the chance to bind to the combining site of the antibody.
10. At the end of the incubation step, transfer the contents of each of the three tubes to clean reaction vessels containing one of the three identical test samples.

For Western blotting strips:

Incubate the strips with the pre-incubated antibody preparations for 1 hour at room temperature or overnight at 4°C .

Wash each strip four times, five minutes each, to remove unbound antibody. Transfer each strip to a new solution containing a labeled secondary antibody [e.g., goat F(ab)₂ anti-rabbit IgG alkaline phosphatase conjugate or goat F(ab)₂ anti-rabbit IgG horseradish peroxidase conjugate].

Remove unbound secondary antibody by thorough washing, and develop the signal using your chemiluminescent reagents and instrumentation.

The signal obtained with antibody incubated with the "Water Only, No Peptide Control" (Tube 1), represents the maximum signal in the assay. This signal should be eliminated by preincubation with the "Phosphopeptide" (Tube 2), while pre-incubation with the "Non-Phosphopeptide" (Tube 3) should not impact the signal. If the "Phosphopeptide" only partially eliminates the signal, repeat the procedure using twice the volume of water or peptide solutions listed in Step 8. If partial competition is seen following pre-incubation with the "Non-Phosphopeptide", repeat the procedure using half the volumes of water or peptide solutions listed in Step 8.

Pictures:
Peptide Competition and Phosphatase

Treatment Extracts of 3T3-L1 cells untreated (1), treated with 100 nM Wortmanin for 90 minutes (2), stimulated with 100 nM Wortmanin for 90 minutes plus 100 nM insulin for 60 minutes (3), stimulated with 100 nM insulin for 20 minutes (4), or stimulated with 100 nM insulin alone for 10 minutes (5-10), were resolved by SDS-PAGE on an 8% Tris-glycine gel and transferred to PVDF. The membrane was left untreated (1-7, 9, 10) or treated with lambda (λ) phosphatase (8), blocked with a 5% BSA-TBST buffer for one hour at room temperature, then incubated with the glycogen synthase [pSpS641/645] antibody for two hours at room temperature in 3% BSA-TBST buffer, following prior incubation with: no peptide (1-4), the non-phosphopeptide corresponding to the phosphopeptide immunogen (5), a generic phosphoserine-containing peptide (6), the phosphopeptide immunogen (7) or the phosphopeptides corresponding to glycogen synthase sites [pSpS8/11] (9) or [pSpS649/653] (10). After washing, the membrane was incubated with goat F(ab')₂ anti-rabbit IgG HRP conjugate and signals were detected using the Pierce SuperSignal™ method. The data show that Wortmanin activates the phosphorylated antibody signal (by inhibiting Akt and thus blocking inhibition of GSK3), while insulin inhibits the signal (by activating Akt and thus inhibiting GSK3). The data also show that only the phosphopeptide corresponding to glycogen synthase [pSpS641/645] blocks the signal and that phosphatase stripping eliminates the signal, verifying that the antibody is indeed phosphorylation site-specific.

