

Monoclonal Antibody to Transforming Growth Factor Alpha

Alternate names:	EGF-like TGF, ETGF, TGF-A, TGF-alpha, TGFA
Catalog No.:	DM3221
Quantity:	0.5 ml
Concentration:	0.3 mg/ml
Host / Isotype:	Mouse / IgG1
Clone:	B326
Immunogen:	Synthetic TNF-alpha 34-43 decapeptide linked carrier.
Applications:	Immunohistochemistry. Western Blotting. This antibody can be used on formalin-fixed, paraffin embedded tissue sections. Prolonged fixation in buffered formalin can destroy the epitope. The antibody may be used at a dilution of with AutoProbe III (Cat. No. 08-803). It is recommended that this product be used on frozen tissue sections or specimens. Other applications not tested. Optimal dilutions of this antibody are dependent on conditions and should be determined by the user. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts with transforming growth factor a, a small 6 kD peptide. It does not cross-react with epidermal growth factor (EGF) or transforming growth factor b (TGF b).
Storage:	Store the antibody at 4°C. Do not freeze! Shelf life: one year from despatch.

Aliquoting Instructions: Do not dilute the entire reconstituted solution at once. Withdraw aliquots as needed with a micropipette and keep concentrated stock at 4°C. Dilute according to the particular application being used. In general, the 0.05M borate pH 8.0 containing 0.15M sodium chloride, 0.02% sodium azide, is a good diluent to use with most antibodies. Avoid diluting the entire contents of the vial at once since the diluted solution