

Polyclonal Antibody to Somatostatin

Catalog No.:	DP3122
Quantity:	0.5 ml
Concentration:	2.5 mg/ml total IgG
Host:	Rabbit
Immunogen:	Synthetic somatostatin
Format:	State: Liquid Purification: Purified Buffer System: 20 mM tris-borate, 150 mM Sodium Chloride, BMC-6 carrier polysaccharides, carrier protein, and 0.05% Sodium Azide, pH 7.5
Applications:	This antibody can be used on formalin-fixed, paraffin-embedded tissue sections (1/50). Prolonged fixation in buffered formalin can destroy the epitope. It is recommended that this product be used on frozen tissue sections or specimens. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts with somatostatin, a growth hormone-release inhibitor. Species: Mammals.
Storage:	Refrigerate at 4°C. Do not

Add. Informations: Aliquoting Instructions: Do not dilute the entire reconstituted solution at once. Withdraw aliquots as needed with a micropipette and keep concentrated stock at 4°C. Dilute according to the particular application being used. Other species not tested.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

