

Polyclonal Antibody to Human TRADD (TNF-R1-Associated Death Domain)

Catalog No.:	DP3136
Quantity:	0.5 mg
Concentration:	1 mg/ml
Host:	Rabbit
Immunogen:	A synthetic peptide from C-terminus of human TRADD.
Applications:	Western Blotting: Ab 5-10µg/ml for 2 hrs at RT. Immunohistology (formalin/paraffin): Use Ab at 10-20µg/ml for 30 min at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. Recommended positive control: Raji cells or exponentially growing any cultured human cells. Other applications not tested. Optimal dilutions of this antibody are dependent on conditions and should be determined by the user. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	TRADD (TNF-R1-Associated Death Domain) is an important mediator of FAS and TNFr1 associated apoptotic signaling. Reacts with HeLa Cells, Tonsil, and Placenta. Cellular Localization: cytoplasmic. Mol. Wt of Antigen: 34kD
Storage:	Store the antibody at 4°C. Do not freeze! Shelf life: one year from despatch.
General Readings:	1. Hsu H, Xiong J, Goeddel D V. (1995) Cell 81:495-504. 2. Baker S J, Reddy E P. (1996) Oncogen

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com