

## CD68, Macrophage-PG-M1 (PG-M1) Antibody

<b>Catalog No.:</b>	EB05298
<b>Quantity:</b>	0.1 mg
<b>Concentration:</b>	0.36 mg/ml
<b>Background:</b>	This antibody detects a 110 kD glycoprotein and recognizes macrophages in a wide variety of human tissues, including Kupffer's cells and macrophages in spleen, gut, lung, and bone marrow.
<b>Host / Isotype:</b>	Goat
<b>Immunogen:</b>	Gaucher's cell
<b>Format:</b>	<b>State:</b> Concentrated <b>Buffer System:</b> 20 mM Tris-Borate, 150 mM Sodium Chloride, dialyzed media RPMI 1640/D-MEM containing Fetal Bovine Serum, BMC-6 Carrier Polysaccharides, Carrier Protein, and 0.05% Sodium Azide, pH 7.5.
<b>Applications:</b>	IHC-Fr, IHC-P, WB Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	Human. Others not tested.
<b>Storage:</b>	Keep as concentrated solution, aliquot and store at 4

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)