



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.incuna.com

sales@everestbiotech.com

sales@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for
diagnostic or therapeutic use.**

Storage: For long-term storage
keep aliquots at -20°C. (Store no
longer than 12 months at 4°C).
Minimize freezing and thawing.

EB06524 - Goat Anti-RXR beta Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: DADB-100D22.1, MHC class I promoter binding protein, retinoid X receptor, beta, H-2RIIBP, MGC1831, RCoR-1, DAUDI6, NR2B2, RXRB

Official Symbol: RXRB

Accession Number(s): NP_068811.1

Human GeneID(s): [6257](#)

Immunogen

Peptide with sequence CRDGMGDSGRDSRSP, from the internal region of the protein sequence according to NP_068811.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 55kDa band observed in lysates of cell line A431 (calculated MW of 56.9kDa according to NP_068811.1). In transfected HEK293 transiently expressing RXRB a band of approx. 60kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow


Background Reference

Samad TA, Krezel W, Chambon P, Borrelli E.


Regulation of dopaminergic pathways by retinoids: activation of the D2 receptor promoter by members of the retinoic acid receptor-retinoid X receptor family.

Proc Natl Acad Sci U S A. 1997 Dec 23;94(26):14349-54.

PMID: 9405615



EB06524 staining (1 μ g/ml) of A431 lysate (RIPA buffer, 35 μ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



HEK293 overexpressing RXRB (RC200516) and probed with EB06524 (mock transfection in first lane), tested by Origene.