

Catalog Number: GT15024 Host: Goat

Product Type: Affinity Purified Goat IgG Species Reactivity: Rat, mouse Immunogen Sequence: Recombinant rat fractalkine Format: Liquid 1mg/ml

0.2 µm filtered solution in phosphate-

buffered saline (PBS) with 5% trehalose.

Applications: Immunohistochemistry: 1-2 µg/mL

Western Blot: 0.1 - 0.2 µg/mL

Only tested for Western using recombinant rat fractalkine

Dilutions listed as a recommendation. Optimal dilution should be determined by investigator.

Storage: The antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Antibody

can be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six

months without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Application Notes

Western Blot: Only tested for Western using recombinant rat Factalkine

Immunohistochemistry: Antiserum was used on fixed tissue.

This antibody can be used with the appropriate secondary reagents to detect Fractalkine on tissues dissected from animals that were fixed by vascular perfusion with 4% paraformaldehyde/PBS (pH 7.4) and followed by perfusion with a 10% sucrose solution in 0.1 M phosphate buffer (pH 7.2).

FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.

V. 01/04