

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 <u>US-info@acris-antibodies.com</u>

## PA1049 OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## **Rat Interferon tau - Purified** IFN tau Alternate names: PA1049 **Catalog No.: Quantity:** 2 µg **Concentration:** 1,0 mg/ml **Background:** IFN-tau is also known as TP-1 (trophoblast protein-1) is a new class of type I IFN that is secreted by the trophoblast and is the signal for maternal recognition of pregnancy in sheep. IFN- tau has potent immunosuppressive and antiviral activities similar to other type I IFN but is less cytotoxic than IFN-alpha and IFN-beta. The current investigation concerns the effect of recombinant ovine IFN- tau (rOIFN- tau) on the modulation of MHC class I and II expression on cloned mouse cerebrovascular endothelial (CVE) cells. IFN-tau induced tyrosine phosphorylation of Stat1 and upregulated the expression of MHC class I on CVE. One proposed action by which type I IFN reduces the relapse rate in MS is via interference with IFN-y-induced MHC class II expression. IFN- tau was shown to downregulate IFN-y-induced MHC class II expression on CVE and, hence, may be of potential therapeutic value in downregulating inflammation in the central nervous system (CNS). IFN- tau did not upregulate the expression of MHC class II on CVE. IFN- tau also inhibited the replication of Theiler's virus in CVE. **Species:** Rat Source: E. coli, Escherichia Coli Format: State: Lyophilized purified protein (sterile filtered) **Purity:** >95% Greater than 95.0% as determined by both: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE. Buffer System: PBS pH-7.4 **Reconstitution:** Restore in sterile 18Mohm-cm H2O not less than $100\mu g/ml$ , which can then be further diluted to other aqueous solutions **Description:** Interferon-Tau Ovine Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 172 amino acids. The IFN-Tau is purified by proprietary chromatographic techniques. AA Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Cys-Tyr-Leu-Ser-Arg. **Specific Activity:** 1 x 10e7 IU/mg Molecular weight: 17 kDa 19914.7 Dalton Add. Information: Protein quantitation was carried out by two independent methods: 1. UV spectroscopy at 280 nm using the absorbency value of 1.027 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.







## PA1049: Rat Interferon tau - Purified

 analysis program of protein sequences (IntelliGenetics).
Analysis by RP-HPLC, using a calibrated solution of IFN-Tau as a Reference Standard.
Storage: Prior to reconstitution stable at RT for 3 weeks.
Following reconstitution store the protein undiluted at 2-8°C for one week or (in aliquots) at -20°C for longer.
Avoid repeated freezing and thawing.
For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Shelf life: one year from despatch.



OG/20121030