

Human Leptin Triple Antagonist - Purified

| | |
|--------------------------|--|
| Catalog No.: | PA1117X |
| Quantity: | 0.1 mg |
| Concentration: | 1,0 mg/ml |
| Background: | Leptin Antagonist Triple Mutant Human Recombinant is a single non-glycosylated polypeptide chain containing 146 amino and additional Ala at N-terminus acids and having a molecular weight of 16 kDa, Leptin was mutated, resulting in L39A/D40A/F41A. Leptin Antagonist Triple Mutant Human Recombinant was purified by proprietary chromatographic techniques. |
| Species: | Human |
| Source: | E. coli, E.coli |
| Format: | State: Lyophilized purified protein (sterile filtered) Purity: >98% Greater than 98.0% as determined by: (a) Gel filtration analysis. (b) Analysis by SDS-PAGE. Buffer System: 0.0045mM NaHCO ₃ Reconstitution: Restore in sterile 0.4% NaHCO ₃ adjusted to pH 8-9, not less than 100µg/ml, which can then be further diluted to other aqueous solutions. |
| Description: | Leptin Antagonist Triple Mutant Human Recombinant is a single non-glycosylated polypeptide chain containing 146 amino and additional Ala at N-terminus acids, Leptin was mutated, resulting in L39A/D40A/F41A. Leptin Antagonist Triple Mutant Human Recombinant was purified by proprietary chromatographic techniques. AA Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Ala-Val-Pro-Ile-Gln. Biological Activity: Leptin triple antagonist is capable of inhibiting leptin-induced proliferation of BAF/3 cells stably transfected with the long form of human leptin receptor. It also inhibits various leptin effects in several in vitro bioassays. Molecular weight: 16 kDa 16 kDa |
| Add. Information: | Protein quantitation was carried out by UV spectroscopy at 280 nm using the absorbency value of 0.88 as the extinction coefficient for a 0.1% (1mg/ml) solution at pH 8.0. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). |

Storage:

Prior to reconstitution stable at RT for several weeks.
Following reconstitution (at > 0.1 mg/ml and up to 2 mM) store the protein undiluted at 2-8°C for one week
or (in aliquots) at -20°C for longer.
Avoid repeated freezing and thawing.
At lower concentration addition of a carrier protein (0.1% HSA or BSA) is suggested.
Shelf life: one year from despatch.