

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## **R1075** Polyclonal Antibody to Cholesterol Oxidase - Serum

Alternate names:	CHOD, Cholesterol oxidase, EC 1.1.3.6
Quantity:	2 ml
Concentration:	90.0 mg/ml (by Refractometry)
Background:	Cholesterol Oxidases exist as both type I and type II oxidases and are implicated in bacterial pathogenesis. In addition, they are important as clinical reagents, potential larvicides, and tools in cell biology.
Host:	Goat
Immunogen:	Cholesterol Oxidase [Microorganism].
Format:	<ul> <li>State: Lyophilized Serum</li> <li>Purification: Prepared from monospecific antiserum by a delipidation and defibrination.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% sodium azide as preservative.</li> <li>Reconstitution: Restore with 2.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoprecipitation and most immunological methods requiring high titer and specificity. This product has been assayed against 1.0 ug of Cholesterol Oxidase [Microorganism] in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Goat) and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:8,000 to 1:40,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against purified and partially purified Cholesterol Oxidase [Microorganism]. Cross reactivity against Cholesterol Oxidase from other sources is unknown.
Storage:	Store vial at 2-8°C prior to restoration. Centrifuge product if not completely clear after standing at room temperature. For extended storage aliquot contents and freeze at -20°C or below. This product is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.