

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## **R1151** Polyclonal Antibody to Maltose Phosphorylase - Serum

Quantitu	
Quantity:	2 ml
Concentration:	90.0 mg/ml (by Refractometry)
Background:	Maltose phosphorylase is a dimeric enzyme that catalyzes the conversion of maltose and inorganic phosphate into beta D glucose 1 phosphate and glucose without requiring any cofactors, such as pyridoxal phosphate. The enzyme is part of operons that are involved in maltose/malto oligosaccharide metabolism.
Host:	Goat
Immunogen:	Maltose Phosphorylase [E.coli].
Format:	<ul> <li>State: Lyophilized purified Ig fraction.</li> <li>Purification: Prepared from monospecific antiserum by a delipidation and defibrination.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% sodium azide as preservative.</li> <li>Reconstitution: Restore with 2.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoprecipitation and most immunological methods requiring high titer and specificity. <u>Recommended Dilutions</u> : This product has been assayed against 1.0 ug of Maltose Phosphorylase [E.coli] in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) and (ABTS (2,2'-azino- bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against purified and partially purified Maltose Phosphorylase [E.coli]. Cross reactivity against Maltose Phosphorylase from other sources is unknown.
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.