

Polyclonal Antibody to Maltose Phosphorylase - HRP

Catalog No.:	R1151HRP
Quantity:	20 mg
Concentration:	10.0 mg/ml (by UV absorbance at 280 nm)
Background:	Maltose phosphorylase is a dimeric enzyme that catalyzes the conversion of maltose and inorganic phosphate into beta D glucose 1 phosphate and glucose without requiring any cofactors, such as pyridoxal phosphate. The enzyme is part of operons that are involved in maltose/malto oligosaccharide metabolism.
Host:	Goat
Immunogen:	Maltose Phosphorylase [E.coli].
Format:	State: Lyophilized purified Ig fraction. Purification: Multistep process. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% Gentamicin Sulfate as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer. Label: HRP – Horseradish Peroxidase Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoperoxidase electron microscopy and Immunohistochemistry as well as other peroxidase-antibody based enzymatic assays. Recommended Dilutions: This product has been assayed against 1.0 mg of Maltose Phosphorylase [E.coli] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:10,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Maltose Phosphorylase [E.coli]. Cross reactivity against Maltose Phosphorylase from other sources is unknown.

- Storage:** Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature.
This antibody is stable for one month at 2-8°C as an undiluted liquid.
Dilute only prior to immediate use.
Avoid repeated freezing and thawing.
Shelf life: One year from despatch.
- Caution:** Do Not Use Sodium Azide as Preservative.
- General Readings:** Farr & Nakane, J. Immunol. Methods 47; 129-144. 1981. (Conjugation)