

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-858-888-7900 Fax: +1-858-888-7904 <u>US-info@acris-antibodies.com</u>

## R1156P OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Polyclonal Antibody to Pyranose Oxidase - Ig Fraction

Catalog No.:	R1156P
Quantity:	50 mg
Concentration:	90.0 mg/ml (by Refractometry)
Host:	Goat
Immunogen:	Pyranose Oxidase [E.coli].
Format:	<ul> <li>State: Lyophilized purified Ig fraction.</li> <li>Purification: Prepared from monospecific antiserum by a delipidation and defibrination.</li> <li>Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% sodium azide as preservative.</li> <li>Reconstitution: Restore with 2.0 ml of deionized water (or equivalent).</li> </ul>
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoprecipitation and most Immunological methods requiring high titer and specificity. <u>Recommendeed dilutions</u> : This product has been assayed against 1.0 ug of Pyranose Oxidase [E.coli] in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:50,000 to 1:400,000 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against purified and partially purified Pyranose Oxidase [E.coli]. Cross reactivity against Pyranose Oxidase from other sources is unknown.
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.



OG/20121030