

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11

info-de@origene.com

R1302T Polyclonal Antibody to Chicken IgG F(ab')2 -TRITC-

Alternate names: Chicken Immunoglobulin G, Chicken Immunoglobulin Y

Quantity: 1.5 mg

Concentration: 1.5 mg/ml (by UV absorbance at 280 nm)

Host: Rabbit

Immunogen: Chicken IgG F(ab')2 fragment.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01%

(w/v) sodium azide as preservative.

Label: TRITC – Tetramethylrhodamine isothiocyanante (Molecular Weight 444 daltons)

Absorption / Emission: 550 nm / 570 nm

Molar Ratio: 3.0 moles TRITC per mole of Rabbit IgG.

Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).

Applications: Suitable for Immunomicroscopy (1/500-1/2,500) and Flow cytometry

(1/2,000-1:10,000) or FACS analysis as well as other antibody based fluorescent

assays requiring lot-to-lot consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Chicken IgG coupled to agarose beads followed by solid phase

adsorption(s) to remove any unwanted reactivities.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-

Rabbit Serum, Chicken IgG, Chicken IgG F(ab')2 and Chicken Serum.

No reaction was observed against Chicken IgG F(c).

Storage: Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and

then aliquot contents and freeze at -20°C or below. Centrifuge product if not

completely clear after standing at room temperature.

This antibody is stable for one month at 2-8°C as an undiluted liquid.

Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.

General Readings: J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)