

R1305AP**Polyclonal Antibody to Bovine IgG (Fc specific) - AP**

Alternate names:	Bovine Immunoglobulin G
Quantity:	1 mg
Concentration:	0.41 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Bovine IgG F(c) fragment.
Format:	State: Liquid (sterile filtered) purified Ig fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.05M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: AP – Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)
Applications:	Suitable for Immunoblotting (Western or Dot blot), ELISA and Immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot consistency. Recommended Dilutions: This product has been assayed against 1.0 ug of Bovine IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,500 of the reconstitution concentration is suggested for this product. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Bovine IgG and Bovine Serum.
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Dilute only prior to immediate use. Shelf life: one year from despatch.
General Readings:	Modified from Avrameas and Ternyrook, Immunochemistry 32; 1175 1971. (Conjugation)