

**R1316P****Polyclonal Antibody to Ferret IgG, IgA, IgM [H&L]**

<b>Alternate names:</b>	Ferret IgA, Ferret IgG, Ferret IgM
<b>Quantity:</b>	1 mg
<b>Concentration:</b>	1.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Goat
<b>Immunogen:</b>	Ferret IgG, IgA and IgM whole molecules.
<b>Format:</b>	<b>State:</b> Liquid (sterile filtered) purified Ig fraction. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0 without preservatives.
<b>Applications:</b>	Immunoprecipitation, Immunodiffusion, Conjugation and most immunological methods requiring lot-to-lot consistency, high titer and specificity. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. This antibody is suitable for the detection of all Ferret immunoglobulin classes, isotypes and chain combinations.
<b>Storage:</b>	Store the antibody (undiluted) at 2-8°C for one month or (in aliquots) at -20°C for longer. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.