

**R1322T****Polyclonal Antibody to Guinea Pig IgG [H&L] -TRITC-**

<b>Alternate names:</b>	Guinea pig Immunoglobulin G
<b>Quantity:</b>	2 mg
<b>Concentration:</b>	2.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Guinea Pig IgG whole molecule.
<b>Format:</b>	<b>State:</b> Lyophilized purified Ig fraction. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 <b>Preservatives:</b> 0.01% (w/v) Sodium Azide <b>Stabilizers:</b> 10 mg/ml BSA (IgG and Protease free) <b>Label:</b> TRITC – Tetramethylrhodamine isothiocyanate (Molecular Weight 444 daltons) <i>Absorption / Emission:</i> 550 nm / 570 nm <i>Molar Ratio:</i> 2.0 moles TRITC per mole of Rabbit IgG. <b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent).
<b>Applications:</b>	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. <u><i>Recommended Dilutions:</i></u> FLISA: 1/10,000-1/50,000. IF Microscopy: 1/1,000-1/5,000. Flow Cytometry: 1/500-1/2,500. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Guinea Pig IgG and Guinea Pig Serum.
<b>Storage:</b>	Store vial at 2-8°C prior to restoration. For extended storage aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
<b>General Readings:</b>	1. J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)