

R1358TR**Polyclonal Antibody to Mouse Lambda (Lambda chain specific)
-Texas Red (™) -**

| | |
|--------------------------|---|
| Quantity: | 1 mg |
| Concentration: | 1.0 mg/ml (by UV absorbance at 280 nm) |
| Host: | Rabbit |
| Immunogen: | Mouse λ (Lambda light chain). |
| Format: | State: Lyophilized purified Ig fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. ; Fluorochrome/Protein Ratio: Label: Texas Red – (™) Sulfonyl Chloride (TR; Molecular Weight 625 daltons) <i>Absorption / Emission:</i> 596 nm / 620 nm <i>Molar Ratio:</i> 2.9 moles Texas Red per mole of Rabbit IgG. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent). |
| Applications: | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. |
| Specificity: | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Mouse IgG and Mouse Serum. Solid phase adsorption(s) against a panel of Mouse IgG kappa, IgM kappa and IgA kappa proteins was performed. Specificity was confirmed by ELISA at less than 1% cross reactivity against other human heavy or light chain isotypes. |
| Storage: | Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch. |
| General Readings: | J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation). |