

**R1433R****F(ab')<sub>2</sub> Fragment of Donkey anti-Sheep IgG (H&L)-PE-**

<b>Alternate names:</b>	Sheep Immunoglobulin G
<b>Quantity:</b>	0.5 mg
<b>Host:</b>	Donkey
<b>Immunogen:</b>	Sheep IgG whole molecule
<b>Format:</b>	<b>State:</b> Lyophilized F(ab') <sub>2</sub> fragments. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride; pH 7.2 with 0.02% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA), IgG and Protease free, as stabilizer. <b>Label:</b> PE – R-Phycoerythrin ( (Molecular Weight 240,000 daltons) from seaweed. <b>Phycobiliprotein Concentration:</b> 0.5 mg/ml (determined by absorbance = 82.0 at 565 nm for a 1% solution for only those molecules to which at least one molecule of active antibody is bound) <i>Absorption / Emission:</i> 490 nm, 545 nm and 565 nm / 580 nm <b>Reconstitution:</b> Restore with 1.0 ml of deionized water (or equivalent).
<b>Applications:</b>	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. <b>Recommended Dilutions:</b> The maximum amount of reagent required to stain 1 x 10 <sup>6</sup> cells in Flow Cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. As a general guideline dilutions of 1:50 to 1:200 should be suitable for most applications. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by Immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Donkey Serum, Sheep IgG and Sheep Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG Fc, or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit or Rat Serum Proteins.
<b>Storage:</b>	Store the antibody (undiluted) at 2-8°C prior to opening. <b>DO NOT FREEZE!</b> Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis. Dilute only prior to immediate use. Shelf life: One year from despatch.