

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

OriGene EU

R1437C2

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Polyclonal Antibody to Hamster IgG [H&L] -Cy2-

Alternate names: Hamster Immunoglobulin G

Catalog No.: R1437C2
Quantity: 1 mg

Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Host: Goat

Immunogen: Hamster IgG, whole molecule.

Format: State: Lyophilized purified Ig fraction.

Purification: Immunoaffinity chromatography.

Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG

and Protease free) as stabilizer.

Label: Cy2 – (Cyanine 2-OSu; Molecular Weight 897 daltons)

Absorption / Emission: 489 nm / 505 nm

Molar Ratio: 2.2 moles Cy2 per mole of Goat IgG (1-4 acceptable). **Reconstitution**: Restore with 1.0 ml of deionized water (or equivalent). (

Applications: Suitable for Immunomicroscopy (1:100-1:1,000) and Flow Cytometry (1:200-1:400) or FACS

analysis as well as other antibody based fluorescent assays requiring lot-to-lot

consistency.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This product was prepared from monospecific antiserum by Immunoaffinity

chromatography using Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities and extensive dialysis against the buffer

stated above.

Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Goat

Serum, Hamster IgG and Hamster Serum.

No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Horse, Human,

Mouse, Rabbit, Rat and Sheep Serum Proteins.

Add. Information: Aggregation Index: (OD463-OD473)/OD490 x 100% = 0.7% (< 4% acceptable)

Store vial at 2-8°C prior to restoration. For extended storage mix with glycerol to 50% and

then aliquot contents and freeze at -20 $^{\circ}$ C or below. Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8 $^{\circ}$ C as

an undiluted liquid. Dilute only prior to immediate use.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

TUV HORD CERT CRIENT OF THE PARK AND THE THE P