



OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200
Rockville, MD 20850
UNITED STATES
Phone: +1-858-888-7900
Fax: +1-858-888-7904
US-info@acris-antibodies.com

RA100D8

OriGene EU

Acris Antibodies GmbH
Schillerstr. 5
32052 Herford
GERMANY
Phone: +49-5221-34606-0
Fax: +49-5221-34606-11
info@acris-antibodies.com

S. aureus Protein A (IRDye800 conjugated) - IRDYE800

Catalog No.:	RA100D8
Quantity:	0.5 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Background:	Protein A is a surface protein of S.aureus which binds IgG molecules by their Fc region. In serum, the bacteria will bind IgG molecules in the wrong orientation on their surface which hinders opsonization and phagocytosis. Mutants of S. aureus lacking protein A are more efficiently phagocytosed in vitro, and mutants in infection models have diminished virulence. Due to its affinity for the Fc region of many mammalian immunoglobulins, protein A is considered a universal reagent in biochemistry and immunology.
Species:	S. aureus
Format:	State: Lyophilized Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml BSA and 0.01% (w/v) Sodium Azide Label: IRDye(TM)800 (MW 1166.2) Fluorochrome/Protein Ratio: 4.0 IRDye(TM)800 per mole of Protein A Absorption/Emission: 778 nm / 806 nm Reconstitution: Restore with 0.5 ml of deionized water (or equivalent).
Applications:	Western Blot: 1/10,000 to 1/25,000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Description:	Single precipitin arc against anti-Protein A.
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the protein undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.