

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11

info-de@origene.com

SM036P Monoclonal Antibody to CD48 - Purified

Alternate names: B-lymphocyte activation marker BLAST-1, BCM1, BCM1 surface antigen, BLAST1,

Leukocyte antigen MEM-102, MEM-102, TCT.1

Quantity: 0.25 mg **Concentration:** 1.0 mg/ml

Background: CD48 is a ligand for the mouse CD2 cell surface antigen.

Uniprot ID: P18181

NCBI: NP 031675.1

GeneID: 12506

Host / Isotype: Rat / IgG2a

Recommended Isotype

SM15P, SM15PX

Controls: Clone:

OX-78

Recombinant soluble Mouse CD48. Immunogen:

Spleen cells from immunised AO Rats were fused with cells from the Rat Y3 myeloma

Format: **State:** Liquid purified IgG fraction from tissue culture supernatant

Purification: Affinity Chromatography on Protein G

Preservatives: 0.09% Sodium Azide

Flow Cytometry: Use 10 μ l of 1/50-1/100 diluted antibody to label 10⁶ cells in 100 μ l. **Applications:**

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

This antibody labels Mouse thymocytes, B and T cells and macrophages. **Specificity:**

Species: Mouse.

Other species not tested.

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

> Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: 1. Kato K, Koyanagi M, Okada H, Takanashi T, Wong YW, Williams AF, et al. CD48 is a

counter-receptor for mouse CD2 and is involved in T cell activation. J Exp Med. 1992

Nov 1;176(5):1241-9. PubMed PMID: 1383383.

2. Lapaque N, Forquet F, de Chastellier C, Mishal Z, Jolly G, Moreno E, et al.

Characterization of Brucella abortus lipopolysaccharide macrodomains as mega rafts.

Cell Microbiol. 2006 Feb;8(2):197-206. PubMed PMID: 16441431.

3. Kashiwakura, Y. et al. (2012) CD2-mediated regulation of peripheral CD4(+) CD25(+) regulatory T cell apoptosis accompanied by downregulation of Bim. Immunology. Dec

24. [Epub ahead of print]

4. Wild MK, Cambiaggi A, Brown MH, Davies EA, Ohno H, Saito T, et al. Dependence of T cell antigen recognition on the dimensions of an accessory receptor-ligand complex.



J Exp Med. 1999 Jul 5;190(1):31-41. PubMed PMID: 10429668.

Pictures:

Staining of Mouse spleen cells with Rat Anti Mouse CD48 Antibody Cat.-No SM036.

