

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

SM088P OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to T Cell Receptor (TCR) Vb 7 -Purified

Alternate names:	TCR V beta-7, TCR Vb7
Catalog No.:	SM088P
Quantity:	0.2 mg
Concentration:	0.2 mg/ml
Host / Isotype:	Rat / IgG2b
Clone:	TR310
Immunogen:	T-cell lymphoma TK1. Spleen cells from immunised mice were fused with cells of the P3-X63 Ag8.3 rat myeloma cell line.
Format:	State: Liquid purified IgG Purification: Affinity chromatography on Protein A Buffer System: PBS containing 0.09% Sodium Azide
Applications:	Flow cytometry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts with the TCR Vb7 bearing T cells. Note that TCR Vb7 may not be present in mouse strains expressing MIS-1a haplotype. Species: Mouse. Other species not tested.
Storage:	Store the antibody at 4-8°C for one month or at -20°C for longer. This product should be stored undiluted. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	 Okada CY, Holzmann B, Guidos C, Palmer E, Weissman IL. Characterization of a rat monoclonal antibody specific for a determinant encoded by the V beta 7 gene segment. Depletion of V beta 7+ T cells in mice with Mls-1a haplotype. J Immunol. 1990 May 1;144(9):3473-7. PubMed PMID: 1691759. Suhihara, S. et al. (1993) Autoimmune thyroiditis induced in mice depleted of particular T cell subsets. J. Immunol. 150: 683-694. Ignatowicz, L. et al. (1994) Identification of two Vb7-specific viral superantigens. J. Immunol. 152: 65-71.



OG/20130615