

SM1097AS**Monoclonal Antibody to CD35 / C3BR - Azide Free**

Alternate names:	C3b/C4b receptor, CR1, Complement receptor type 1
Quantity:	50 µg
Concentration:	1 mg/ml
Background:	CD35 is a membrane glycoprotein found on peripheral blood cells, glomerular podocytes, and follicular dendritic cells. The protein is a receptor for complement components C3b and C4b and regulates the activity of the complement cascade. Variation in this protein is the basis of the Knops blood group system. The two most common alleles, F and S, differ by 8 exons and are thought to be the result of an unequal crossover event. A secreted form of the protein present in plasma has been described, but its full length nature has not been determined.
Uniprot ID:	P17927
NCBI:	NP_000564
GeneID:	1378
Host / Isotype:	Mouse / IgG1
Clone:	E11
Immunogen:	Human acute monocytic leukaemia cells. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Format:	State: Liquid purified IgG fraction from Tissue Culture Supernatant Purification: Affinity Chromatography on Protein A Buffer System: PBS, pH 7.4 Preservatives: None Stabilizers: None
Applications:	Flow Cytometry: 1/50 - 1/100. Immunohistochemistry on Frozen Sections. Immunohistochemistry on Paraffin Sections: 10 µg/ml. Western Blot. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody is a single chain cell surface glycoprotein which exists in four allotypic forms (A,B,C,D) of 190kD, 220kD, 160kD and 250kD respectively. The molecule is expressed by granulocytes, monocytes, B cells and some T cells. Functionally CD35 acts as a receptor for the C3b component of complement. Species: Human, Cynomolgus monkey, Baboon, Rhesus Monkey. Other species not tested.
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings:

1. Hogg N, Ross GD, Jones DB, Slusarenko M, Walport MJ, Lachmann PJ. Identification of an anti-monocyte monoclonal antibody that is specific for membrane complement receptor type one (CR1). *Eur J Immunol.* 1984 Mar;14(3):236-43. PubMed PMID: 6368248.
2. Yoshino N, Ami Y, Terao K, Tashiro F, Honda M. Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. *Exp Anim.* 2000 Apr;49(2):97-110. PubMed PMID: 10889948.
3. Birmingham, D. J. et al. (1996) The baboon erythrocyte complement receptor is glycoposphatidyl/insitolinked protein encoded by homologue of the human CR1 - like genetic element. *J. Immunol.* 157: 2586 - 2592.
4. Sopper S, Stahl-Hennig C, Demuth M, Johnston IC, Dörries R, ter Meulen V. Lymphocyte subsets and expression of differentiation markers in blood and lymphoid organs of rhesus monkeys. *Cytometry.* 1997 Dec 1;29(4):351-62. PubMed PMID: 9415418.

Pictures:

Formalin-Fixed, Paraffin-Embedded Human Kidney stained with Azide Free CR1 Antibody Cat.-No SM1097A (Clone E11) 10 µg/ml after heat-induced antigen retrieval.

