

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

SM2032R Monoclonal Antibody to MHC Class II DQ/DR (polymorphic) - PE

Quantity: 100 Tests

Background: Ovine MHC class II antigens are expressed on several cell types, including B cells,

activated T cells, monocytes, macrophages and dendritic cells.

Host / Isotype: Mouse / IgG1

Clone: 28.1

Immunogen: Ovine alveolar macrophages. Spleen cells from immunised BALB/c mice were fused

with cells of the mouse NS1 myeloma cell line.

Format: State: Lyophilized purified IgG

Purification: Affinity chromatography on Protein G

Buffer System: PBS, pH 7.4 containing 0.09% Sodium Azide and 1% Bovine Serum

Albumin

Label: PE – R. Phycoerythrin (RPE)

Reconstitution: Restore with 1 ml distilled water

Applications: Flow Cytometry: Neat - 1/10; Use 10µl of the suggested working dilution to label 10e6

cells in 100µl.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: This antibody recognises a polymorphic epitope on MHC class II DQ and DR

molecules. In recent work, this clone was found to recognise ovine MHC II

transfectants DQ - T28.1, DQ - T26.2 and DR - T31.3 but not DR - T8.1. (Ballingall, K. et

al. 1995).

Species: Sheep.

Other species not tested.

Storage: Prior to and following reconstitution store the antibody at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.

General Readings: 1. Puri, N. et al. (1985) Sheep lymphocyte antigens (OLA) II. Major histocompatibility

complex class II molecules.Immunology. 56: 725 - 733.

2. Puri NK, Gogolin-Ewens KJ, Brandon MR. Monoclonal antibodies to sheep MHC class I and class II molecules: biochemical characterization of three class I gene products and four distinct subpopulations of class II molecules. Vet Immunol

Immunopathol. 1987 May;15(1-2):59-86. PubMed PMID: 3303652.

3. Puri NK, Brandon MR. Sheep MHC class II molecules. II. Identification and characterization of four distinct subsets of sheep MHC class II molecules.

Immunology. 1987 Dec;62(4):575-80. PubMed PMID: 3480873.

4. Puri, N. et al. (1987) Sheep MHC class II molecules II. Identification and characterization of sheep MHC class II molecules. Immunology. 62: 575 - 580. 5. Puri NK, de Kretser T, Brandon MR. Monoclonal antibodies to sheep MHC class II

molecules recognize all HLA-D or subsets of HLA-D region products. Hum Immunol.



1987 Nov;20(3):195-207. PubMed PMID: 3501780.

6. Sainte-Marie, G. et al. (1962) A paraffin embedding technique for studies employing immunofluorescence. J. Histochem. Cytochem. 10: 250

7. Ballingall KT, Dutia BM, Hopkins J, Wright H. Analysis of the fine specificities of sheep major histocompatibility complex class II-specific monoclonal antibodies using mouse L-cell transfectants. Anim Genet. 1995 Apr;26(2):79-84. PubMed PMID: 7733511.

Pictures:

Staining of sheep peripheral blood lymphocytes with MOUSE ANTI OVINE MHC CLASS II DQ/DR:RPE (SM2032R).

