

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

SM2091R Monoclonal Antibody to NKG2A/C/E - PE

Alternate names:	CD159a, CD159c, CD159e, NKG2 lectin-like family
Quantity:	100 Tests
Concentration:	0.1 mg/ml
Background:	In mice, NKG2 subunits associate with CD94 to form heterodimers at the surface of natural killer (NK) cells. The CD94/NKG2 heterodimer is the receptor for a non-classical MHC class I ligand, which is Qa-1 in the mouse.
Host / Isotype:	Rat / IgG2a
Clone:	20d5
Immunogen:	CHO transfected cells expressing the B6 allele of NKG2A. Spleen cells from immunised Lewis rats were fused with cells of the mouse P3X63-Ag8.653 myeloma cell line.
Format:	 State: Lyophilized purified IgG fraction. Purification: Affinity Chromatography on Protein G. Buffer System: PBS, pH7.4 containing 0.09% Sodium Azide and 1% Bovine Serum Albumin. Label: PE – R. Phycoerythrin (RPE) Reconstitution: Restore with 1 ml distilled water
Applications:	Flow Cytometry: Neat - 1/2; Use 10µl of the suggested working dilution to label 10e6 cells in 100µl; The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises NKG2A, NKG2C and NKG2E, which are isoforms of the NKG2 lectin-like family. Clone 20d5 is reported to block ligand binding to the receptor. We recommend the use of SM2091LE for this purpose. Species: Mouse. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.
General Readings:	 Vance RE, Jamieson AM, Raulet DH. Recognition of the class Ib molecule Qa-1(b) by putative activating receptors CD94/NKG2C and CD94/NKG2E on mouse natural killer cells. J Exp Med. 1999 Dec 20;190(12):1801-12. PubMed PMID: 10601355. Vance RE, Jamieson AM, Cado D, Raulet DH. Implications of CD94 deficiency and monoallelic NKG2A expression for natural killer cell development and repertoire formation. Proc Natl Acad Sci U S A. 2002 Jan 22;99(2):868-73. Epub 2002 Jan 8. PubMed PMID: 11782535.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

1/1